



THE ROYAL INSTITUTE OF TECHNOLOGY
STOCKHOLM
SWEDEN



DEPARTMENT OF
INFORMATION PROCESSING
COMPUTER SCIENCE

REPORT FROM
SYMPOSIUM ON "NEW APPLICATIONS AND
NUMERICAL METHODS FOR DIFFERENTIAL
EQUATIONS AND SIMILAR PROBLEMS"
APRIL 17-19, 1974

TRITA-NA-7408



ABSTRACT

...way coupling between the nonlinear I.V.P. (1) and this ...
A clear recognition of the underlying methodology is a prerequisite for the
proper estimation of parameters in scientific models. The notions of consis-
tency and uniqueness can be instrumental in this respect. When experimental
data are analysed, in advance a model of the system under investigation has
to be known only with a finite number of degrees of freedom. Now a model is
called consistent with respect to a set of experimental data if no discrepancy
exists between data and model. In general a model will be consistent for
parameter values from a subset of the a priori allowed parameter set. A
model is unique if all parameters are determined by the data, such that there
is no freedom left in the model. It is fundamental that one never can prove
a model to be the correct one. However there is the possibility to reject
the wrong ones: from a correct operational model (i.e. a model which is
consistent and unique) it should be possible to predict future behaviour.
Specially, in this light it is clear that not only estimates of the para-
meters have to be determined, but also estimates of their reliability.

These ... together with the vector of residuals $Y(p)$, enable
The estimation of a set of parameters is reduced to the solution of a
nonlinear least squares problem as follows. Let be given a set of
experimental data $\{(t_i, y_i)\}$ - y_i represents the result of a measurement on
(some component of) y at time t_i , and let the model be given by the initial
value problem

$$\begin{aligned} y'(t,p) &= f(t,y,p) , & (1) \\ y(0,p) &= y_0(p) . \end{aligned}$$

As an estimate of the parameter (vector) p is considered that value of p
which minimizes

$$S(p) = \sum (y(t_i,p) - y_i)^2 = \|Y(p)\|^2 .$$

The minimization is performed by Marquardt's method.

This method requires values of $\partial y(t,p)/\partial p$, which are obtained by solving
the additional set of differential equations

$$\begin{aligned} y'_p(t,p) &= f_p(t,y,p) + f_y(t,y,p) \cdot y_p(t,p), & (2) \\ y'_p(0,p) &= \partial y_0(p)/\partial p . \end{aligned}$$

The one-way coupling between the nonlinear I.V.P. (1) and this linear set of differential equations (2) is exploited by the use of implicit linear multistep methods (Adams-Moulton or Gear's stiffly stable methods, cf. [1]).

Marquardt's method is realized by means of the singular value decomposition of the Jacobian matrix $J = \partial y(t_i, p) / \partial p$; [cf. 2,3]. With $J = U \Lambda V^T$ the iteration process becomes

$$p^{(n+1)} = p^{(n)} - V(\Lambda^2 + \lambda)^{-1} \Lambda U^T Y(p^{(n)})$$

with some $\lambda = \lambda^{(n)} > 0$.

Initial estimates can be obtained in the usual way (i.e. from steady states or from raw estimates of derivatives) or by combining the parameter estimation process with multiple shooting.

The singular values Λ and the corresponding columnvectors of V (eigenvectors of $J^T J$) yield important information about correlations between the different parameters. These matrices, together with the vector of residuals $Y(p)$, enable us to a certain extent to judge of the consistency, uniqueness and accuracy of the parameters determined.

A program for the determination of parameters in initial value problems is described in [1]. A revised version will become available in [2] and will be described in detail by Van Domselaar [4].

References

- [1] P.W. Hemker. Parameter estimation in nonlinear differential equations. Mathematical Centre report MR 134/72.
- [2] NUMAL, a library of numerical procedures in ALGOL 60. (P.W. Hemker, W. Hoffmann, P.J. v.d. Houwen and N. Temme eds.) Mathematical Centre, Amsterdam.
- [3] J.C.P. Bus, B. v. Domselaar, J. Kok. Nonlinear least squares estimations. Mathematical Centre (to appear).
- [4] B. v. Domselaar. Report on nonlinear parameter estimation in differential equations. Mathematical Centre, Amsterdam (to appear).